

GLUCAGEN EMERGENCY KIT (GLUCAGON EMERGENCY KIT)

LOW BLOOD SUGAR CAN HAPPEN WHEN A PERSON TAKES TOO MUCH INSULIN, DOESN'T EAT ON TIME OR DOESN'T EAT ENOUGH, OR EXERCISES TOO MUCH.

WHEN ACTION IS TAKEN QUICKLY, LOW BLOOD SUGAR CAN BE TREATED WITH GLUCOSE TABLETS, A SUGARY SOFT DRINK OR JUICE.

HOWEVER, IF ACTION IS NOT TAKEN QUICKLY, OR A PERSON CANNOT FEEL SYMPTOMS OF LOW BLOOD SUGAR, THE PERSON COULD PASS OUT AND REQUIRE HELP FROM ANOTHER PERSON.

NEVER PUT ANYTHING IN THE MOUTH OF AN UNCONSCIOUS PERSON. THIS IS DANGEROUS AND COULD MAKE THEM CHOKE.

THIS IS WHERE YOUR EMERGENCY KIT COMES IN. FOLLOW THESE SIMPLE STEPS:

- ***INSERT THE NEEDLE THROUGH THE RUBBER STOPPER ON THE GLUCAGON VIAL. INJECT ALL LIQUID IN THE SYRINGE INTO THE RUBBER STOPPER.***
- ***LEAVE THE SYRINGE IN PLACE AND GENTLY SHAKE THE VIAL UNTIL THE POWDER IS MIXED AND THE LIQUID IS CLEAR.***
- ***WITH THE NEEDLE STILL IN THE VIAL, TURN THE VIAL UPSIDE DOWN (KEEP THE NEEDLE IN THE LIQUID) AND SLOWLY WITHDRAW ALL THE LIQUID INTO THE SYRINGE***
- ***INSERT THE NEEDLE INTO THE TOP OF THE LEG (AS DEMONSTRATED BY NURSE), AND INJECT ALL OF THE LIQUID.***
- ***TURN THE PERSON ON HIS/HER SIDE AND PROP WITH PILLOWS IN CASE OF NAUSEA.***
- ***HE/SHE SHOULD AWAKEN WITHIN 15 MIN. IF NOT, CALL 911. AS SOON AS PATIENT IS AWAKE AND ABLE TO SWALLOW, GIVE THEM A SUGARY DRINK OR JUICE AND SOMETHING TO EAT, LIKE PEANUT BUTTER OR CHEESE AND CRACKERS OR A MEAT SANDWICH.***